

Josephson Effect between Condensates with Different Internal Structures

Munehiro Nishida,^{1,2} Noriyuki Hatakenaka,^{1,3} and Susumu Kurihara²

¹Low Temperature Laboratory, Helsinki University of Technology, P.O. Box 2200, FIN-02015 HUT, Finland

²Department of Physics, Waseda University, Okubo, Shinjuku-ku, Tokyo 169-8555, Japan

³NTT Basic Research Laboratories, Atsugi, Kanagawa 243-0198, Japan

(Dated: February 1, 2008)

A general formula for Josephson current in a wide class of hybrid junctions between different internal structures is derived on the basis of the Andreev picture. The formula extends existing formulae and also enables us to analyze novel *B*-phase/*A*-phase/*B*-phase (*BAB*) junctions in superfluid ³He systems, which are accessible to experiments. It is predicted that *BAB* junctions will exhibit two types of current-phase relations associated with different internal symmetries. A “pseudo-magnetic interface effect” inherent in the system is also revealed.

PACS numbers: 67.57.-z, 74.50.+r, 74.80.Fp, 67.57.Fg

Superfluid systems with internal degrees of freedom produce diverse ordered structures, which provide a new arena for exploring the fertile physics behind them. The Josephson effect [1] extracts the global phase resulting from the spontaneous breakdown of gauge symmetry. Moreover, it also provides us with information on the internal structures formed from other broken symmetries since quasiparticles acquire an excess phase from the internal structures while traveling through the junction. In fact, so-called π junctions associated with high- T_c superconductors provide convincing evidence for *d*-wave symmetry [2, 3], and recent experiments on Pb-Sr₂RuO₄-Pb Josephson junctions [4] could be explained by *p*-wave symmetry in Sr₂RuO₄ [5, 6]. Moreover, metastable π states observed in superfluid ³He weak links [7, 8, 9] are considered to be a signature of texture due to the internal degrees of freedom of *p*-wave order parameters [10, 11]. In the above ways, order parameter configurations have been investigated by using Josephson current-phase relations.

Since a wide variety of structures could be realized in systems with internal degrees of freedom such as the textures in superfluid ³He systems, we require a general theory for the Josephson effect to enable us to identify the structures. In this Letter, we derive a general formula for Josephson current, applicable to a wide class of hybrid junctions between different internal structures in unitary states, on the basis of the Andreev picture [12] of the Josephson effect developed by Furusaki and Tsukada (FT) [13], including explicit expressions for Andreev reflection coefficients reflecting their structures.

Consider a junction composed of three different superfluid regions separated with flat interfaces perpendicular to the *z*-axis. The interfaces are located at $z = 0$ and $z = L$. We shall denote these three regions by L ($z < 0$), C ($0 < z < L$), and R ($L < z$), respectively. We assume that the order parameter is uniform in each region. We also assume that the effective mass m , Fermi velocity v_F , and Fermi wave number k_F are the same in all regions. The potential barriers at the interfaces are ignored for

simplicity. We shall use a triad $(\hat{x}, \hat{y}, \hat{z})$ as the base for a frame of reference.

The dc Josephson current I can be calculated from the temperature Green's function, $\mathcal{G}(z, z', \hat{\mathbf{k}}_{\parallel}, \omega_n)$, which is obtained by an analytic continuation, $E + i0^+ \rightarrow i\omega_n$, from the retarded Green's function constructed by scattering data. Here ω_n is the Matsubara frequency defined by $\omega_n = \pi(2n + 1)/\beta$ ($n = 0, \pm 1, \pm 2, \dots$) at temperature $T = 1/k_B\beta$ and $\hat{\mathbf{k}}_{\parallel} \equiv (\hat{k}_x, \hat{k}_y, 0)$ is defined from $\hat{\mathbf{k}} = (\hat{k}_x, \hat{k}_y, \hat{k}_z)$ ($|\hat{\mathbf{k}}| = 1$) along the classical trajectory. Note that the Green's functions are extended in the form of a 4×4 matrix due to the spin and particle-hole degrees of freedom required for triplet pairing. Taking the particle number conservation into account, the current at $z = 0$ is calculated by

$$I = \frac{\hbar}{4im\beta} \lim_{z, z' \rightarrow 0} \left(\frac{\partial}{\partial z} - \frac{\partial}{\partial z'} \right) \sum_{\omega_n, \hat{\mathbf{k}}_{\parallel}} \text{Tr} \mathcal{G}(z, z', \hat{\mathbf{k}}_{\parallel}, \omega_n). \quad (1)$$

Let us calculate the scattering data necessary for constructing the temperature Green's function. In a quasi-classical approximation, the motion of a quasiparticle of energy E is described by Andreev equations applicable to an arbitrary type of pairing [14] as

$$\begin{pmatrix} (-i\hbar v_F \hat{k}_z \frac{\partial}{\partial z} - E) \hat{\sigma}_0 & \hat{\Delta}^{\hat{\mathbf{k}}} \\ \hat{\Delta}^{\hat{\mathbf{k}\dagger}} & (i\hbar v_F \hat{k}_z \frac{\partial}{\partial z} - E) \hat{\sigma}_0 \end{pmatrix} \begin{pmatrix} u \\ v \end{pmatrix} = 0, \quad (2)$$

where $u = (u_{\uparrow} \ u_{\downarrow})^t$ and $v = (v_{\uparrow} \ v_{\downarrow})^t$. $\hat{\sigma}_0$ is the 2×2 unit matrix. The gap matrix in spin space is denoted by $\hat{\Delta}^{\hat{\mathbf{k}}}$. In each region ($\alpha = \text{L, C, R}$) the gap matrix, $\hat{\Delta}_{\alpha}^{\hat{\mathbf{k}}}$, is assumed to be constant and unitary: $\hat{\Delta}_{\alpha}^{\hat{\mathbf{k}}} \cdot \hat{\Delta}_{\alpha}^{\hat{\mathbf{k}\dagger}} = |\hat{\Delta}_{\alpha}^{\hat{\mathbf{k}}}|^2 \hat{\sigma}_0$. The solutions of the Andreev equations in each region are expressed as

$$\begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} (E + \Omega_{\alpha}^{\hat{\mathbf{k}}}) \hat{\sigma}_0 e^{i\nu_{\alpha} z} & \hat{\Delta}_{\alpha}^{\hat{\mathbf{k}}} e^{-i\nu_{\alpha} z} \\ \hat{\Delta}_{\alpha}^{\hat{\mathbf{k}\dagger}} e^{i\nu_{\alpha} z} & (E + \Omega_{\alpha}^{\hat{\mathbf{k}}}) \hat{\sigma}_0 e^{-i\nu_{\alpha} z} \end{pmatrix} \begin{pmatrix} S_p \\ S_h \end{pmatrix} \\ \equiv \mathbf{M}_{\alpha}^{\hat{\mathbf{k}}}(z) \mathcal{S}, \quad (3)$$

where $\Omega_\alpha^{\hat{\mathbf{k}}} = \sqrt{E^2 - |\Delta_\alpha^{\hat{\mathbf{k}}}|^2}$, $S_p = (S_{p\uparrow} \ S_{p\downarrow})^t$, and $S_h = (S_{h\uparrow} \ S_{h\downarrow})^t$. $S_{p(h)\sigma}$ with $\sigma = \uparrow$ or $\downarrow$ is the probability amplitude for the mode propagating as a particle-like (hole-like) quasiparticle with spin σ . The two amplitudes S_L and S_R are connected by the continuity equation:

$$S_L = M_L^{\hat{\mathbf{k}}}(0)^{-1} M_C^{\hat{\mathbf{k}}}(0) M_C^{\hat{\mathbf{k}}}(L)^{-1} M_R^{\hat{\mathbf{k}}}(L) S_R. \quad (4)$$

Here we introduce scattering coefficient matrices in the following form to express the scattering processes concisely;

$$\hat{s} = \begin{pmatrix} s_{\uparrow\uparrow} & s_{\uparrow\downarrow} \\ s_{\downarrow\uparrow} & s_{\downarrow\downarrow} \end{pmatrix}, \quad (5)$$

where $s_{\sigma\sigma'}$ is the scattering coefficient for the process in which an injected quasiparticle with spin σ' is scattered as a quasiparticle with spin σ . The retarded Green's

function is constructed from the scattering coefficient matrices under the FT prescriptions [13]. For $z \leq z' < 0$, the retarded Green's function is given by

$$\begin{aligned} \mathbf{G}(z, z', \hat{\mathbf{k}}_{\parallel}, E) = & -i \frac{m}{\hbar^2 k_F \hat{k}_z} \\ & \times \left[\frac{e^{-ik_F \hat{k}_z (z-z')} M_L^{\hat{\mathbf{k}}-}(z) \begin{pmatrix} \hat{\sigma}_0 & \hat{a}_2 \\ \hat{0} & \hat{0} \end{pmatrix} M_L^{\hat{\mathbf{k}}-}(z')^\dagger}{(E + \Omega_L^{\hat{\mathbf{k}}-})^2 - |\Delta_L^{\hat{\mathbf{k}}-}|^2} \right. \\ & \left. + \frac{e^{ik_F \hat{k}_z (z-z')} M_L^{\hat{\mathbf{k}}}(z) \begin{pmatrix} \hat{0} & \hat{0} \\ \hat{a}_1 & \hat{\sigma}_0 \end{pmatrix} M_L^{\hat{\mathbf{k}}}(z')^\dagger}{(E + \Omega_L^{\hat{\mathbf{k}}})^2 - |\Delta_L^{\hat{\mathbf{k}}}|^2} \right], \quad (6) \end{aligned}$$

where $\hat{0}$ is the 2×2 zero matrix and $\hat{\mathbf{k}}_- = \hat{\mathbf{k}}_{\parallel} - \hat{k}_z \hat{\mathbf{z}}$. The Andreev reflection coefficients for a process in which a particle-like quasiparticle injected from L is reflected as a hole-like quasiparticle is obtained by solving Eq. (4) as

$$\begin{aligned} \hat{a}_1 = & \left[e^{-i\nu_c L} \left(\hat{\Gamma}_C^{\hat{\mathbf{k}}\dagger} - \hat{\Gamma}_L^{\hat{\mathbf{k}}\dagger} \right) \left(1 - \hat{\Gamma}_C^{\hat{\mathbf{k}}} \hat{\Gamma}_R^{\hat{\mathbf{k}}\dagger} \right) + e^{i\nu_c L} \left(1 - \hat{\Gamma}_L^{\hat{\mathbf{k}}\dagger} \hat{\Gamma}_C^{\hat{\mathbf{k}}} \right) \left(\hat{\Gamma}_R^{\hat{\mathbf{k}}\dagger} - \hat{\Gamma}_C^{\hat{\mathbf{k}}\dagger} \right) \right] \\ & \times \left[e^{-i\nu_c L} \left(1 - \hat{\Gamma}_L^{\hat{\mathbf{k}}} \hat{\Gamma}_C^{\hat{\mathbf{k}}\dagger} \right) \left(1 - \hat{\Gamma}_C^{\hat{\mathbf{k}}} \hat{\Gamma}_R^{\hat{\mathbf{k}}\dagger} \right) + e^{i\nu_c L} \left(\hat{\Gamma}_C^{\hat{\mathbf{k}}} - \hat{\Gamma}_L^{\hat{\mathbf{k}}} \right) \left(\hat{\Gamma}_R^{\hat{\mathbf{k}}\dagger} - \hat{\Gamma}_C^{\hat{\mathbf{k}}\dagger} \right) \right]^{-1}, \quad (7) \end{aligned}$$

where $\hat{\Gamma}_\alpha^{\hat{\mathbf{k}}} = \hat{\Delta}_\alpha^{\hat{\mathbf{k}}}/(E + \Omega_\alpha^{\hat{\mathbf{k}}})$ and $\nu_c = \Omega_C^{\hat{\mathbf{k}}}/\hbar v_F \hat{k}_z$. The matrix $\hat{a}_2$ describes the reverse process. The matrices $\hat{a}_2$ and $\hat{a}_1$ are related by $\hat{a}_2(E, \hat{\mathbf{k}}) = \hat{a}_1(E, \hat{\mathbf{k}}_-)^\dagger$. Substi-

tuting the temperature Green's function obtained by the analytic continuation into Eq. (1), a general formula for the Josephson current at temperature T is derived as

$$I = -\frac{k_F^2}{4\hbar} \int \frac{d^2 \hat{k}_{\parallel}}{(2\pi)^2} \frac{1}{\beta} \sum_{\omega_n} \text{Tr} \left[\frac{\hat{\Delta}_L^{\hat{\mathbf{k}}} \hat{a}_1(\omega_n) + \hat{a}_1(\omega_n) \hat{\Delta}_L^{\hat{\mathbf{k}}}}{\tilde{\Omega}_{nL}^{\hat{\mathbf{k}}}} - \frac{\hat{a}_2(\omega_n) \hat{\Delta}_L^{\hat{\mathbf{k}}-\dagger} + \hat{\Delta}_L^{\hat{\mathbf{k}}-\dagger} \hat{a}_2(\omega_n)}{\tilde{\Omega}_{nL}^{\hat{\mathbf{k}}-}} \right], \quad (8)$$

where $\tilde{\Omega}_{nL}^{\hat{\mathbf{k}}} = \sqrt{\omega_n^2 + |\Delta_L^{\hat{\mathbf{k}}}|^2}$. This is the central result of this Letter. This formula is applicable to any type of Josephson junction between unitary states with any symmetry. Note that our formula still preserves the original FT form expressed by the difference between Andreev reflection coefficients, describing the net current carried by the two processes: the scattering of a particle-like quasiparticle into a hole-like quasiparticle and its reverse process. This formula can indeed be reduced to the FT formula by substituting the s -wave order parameters. Our formula is, therefore, a natural extension of the FT formula.

Let us consider certain special cases reported previously by several authors. Substituting the order parameters used in Ref. 10 into our formulae (7) and (8), and

adopting the limit $L \rightarrow 0$, we obtained an analytic expression for the Josephson current through a pinhole between two $^3\text{He-}B$ reservoirs (Eq. (2) of Ref. 10). This reduces to the Kurkijärvi formula [15] for the B phase if the $\hat{\mathbf{n}}$ vectors of two B -phase reservoirs are parallel. This is the same as the expression for the s -wave supercurrent through a short orifice [16]. By employing the A -phase rather than the B -phase order parameter, the Kurkijärvi formula for the A phase [15] is reproduced. Let us take another limit $\hat{\Delta}_C \rightarrow 0$ with a finite L . This reproduces the current-phase relation of the superconductor/normal-metal/superconductor (SNS) junctions without a potential barrier [17]. The Tanaka-Kashiwaya formula [18] for unconventional *singlet* superconductors without a potential barrier is also reproduced. In addition, the Yamashiro-Tanaka-Kashiwaya results [5] are reproduced

when we employ their order parameters. Thus our formula covers the previous formulae for concrete examples.

Furthermore, our formula enables us to analyze new kinds of junction. One example is *B*-phase/*A*-phase/*B*-phase (*BAB*) junctions in superfluid ^3He systems. This type of junction could be realized by using the experimental setup used to study the Andreev reflection [19] at *AB* interfaces. It is also suggested that a pseudo-*A* phase could be formed around the orifice in ^3He -*B* weak links [20]. In spin-*triplet* Cooper-pair condensates, the boundary condition should be carefully considered to include spin-orbit degrees of freedom. The order parameter of the *A* phase is defined by a triad $(\hat{\mathbf{w}}_1, \hat{\mathbf{w}}_2, \hat{\mathbf{l}})$ in orbital space and a vector $\hat{\mathbf{d}}$ in spin space. For the *B* phase we need a rotational matrix $R(\hat{\mathbf{n}}, \theta)$ with rotational axis $\hat{\mathbf{n}}$ and rotational angle θ relating the spin space to the orbital space. We take the phase angle for the *A* phase to be zero. For an *AB* interface with dipole energy [21, 22], the boundary conditions are $\hat{\mathbf{l}} \parallel \hat{\mathbf{d}} \perp \hat{\mathbf{z}}$, $\hat{\mathbf{d}} = R(\hat{\mathbf{n}}, \theta_L)\hat{\mathbf{w}}_1$, and $\hat{\mathbf{w}}_1 \parallel \hat{\mathbf{z}}$, where $\theta_L = \cos^{-1}(-1/4)$ (Leggett angle). Since the *A* phase always has the ambiguity of $\hat{\mathbf{d}} \rightarrow -\hat{\mathbf{d}}$, $\hat{\mathbf{w}}_1 \rightarrow -\hat{\mathbf{w}}_1$, and $\hat{\mathbf{w}}_2 \rightarrow -\hat{\mathbf{w}}_2$, we choose $\hat{\mathbf{w}}_1 = \hat{\mathbf{z}}$, $\hat{\mathbf{w}}_2 = \hat{\mathbf{x}}$, and $\hat{\mathbf{l}} = \hat{\mathbf{y}}$ without loss of generality. The vector $\hat{\mathbf{n}}$ in the *B* phase is fixed in one of four directions, *i.e.*, $(-\sqrt{3/5}, \pm\sqrt{1/5}, \pm\sqrt{1/5})$ for $\hat{\mathbf{d}} = \hat{\mathbf{l}}$ and $(\sqrt{3/5}, \mp\sqrt{1/5}, \pm\sqrt{1/5})$ for $\hat{\mathbf{d}} = -\hat{\mathbf{l}}$. We use the letters **a**, **b**, **c**, and **d** in turn to specify the different orientations of $\hat{\mathbf{n}}$ for brevity [10]. The order pairs of the letters **aa**, **ab**, etc., thus denote the order parameter configurations at the two interfaces. The order parameters for the configurations of **aa** and **ab**, which become important in the following discussions, are given by

$$\begin{aligned} \hat{\Delta}_L^{\mathbf{k}} &= \Delta_B e^{-i\varphi/2}/2 \\ &\times \left\{ 2i\hat{\sigma}_0\hat{k}_z + \hat{\sigma}_1(\sqrt{3}k_x - k_y) - \hat{\sigma}_3(k_x + \sqrt{3}k_y) \right\} \\ \hat{\Delta}_C^{\mathbf{k}} &= \Delta_A(k_x - i\hat{k}_z)\hat{\sigma}_0 \\ \hat{\Delta}_R^{\mathbf{k}} &= \Delta_B e^{i\varphi/2}/2 \\ &\times \left\{ 2i\hat{\sigma}_0\hat{k}_z + \hat{\sigma}_1(\pm\sqrt{3}k_x - k_y) - \hat{\sigma}_3(k_x \pm \sqrt{3}k_y) \right\}, \quad (9) \end{aligned}$$

where φ is the phase difference and $(\hat{\sigma}_1, \hat{\sigma}_2, \hat{\sigma}_3)$ are Pauli's matrices. Δ_A and Δ_B are the gaps for the *A* phase and the *B* phase, respectively. A rotation of π around the *y*-axis affects the transformations **a** $\leftrightarrow$ **d**, **b** $\leftrightarrow$ **c**, **L** $\leftrightarrow$ **R**, and $\hat{\mathbf{d}} \rightarrow -\hat{\mathbf{d}}$, where we used the ambiguity of the *A* phase. In addition, **aa** and **bb** configurations give the identical current, same as **ab** and **ba**. As a result, we predict two types of current-phase characteristics: type (a) **{aa,bb,cc,dd}** and type (b) **{ab,ba,cd,dc}**. Figure 1 shows the temperature dependence of the Josephson current in (a) **aa** and (b) **ab** configurations for $L = 2\xi_0$ where the coherence length ξ_0 in the *B* phase at zero temperature is given by $\xi_0 = [7\zeta(3)/48]^{1/2}(\hbar v_F/\pi k_B T_c)$ with ζ being the Riemann zeta-function. Here we assume




FIG. 1: Josephson current-phase relations of *BAB* junctions for $L = 2\xi_0$ at temperatures $0.1T_c$, $0.3T_c$, $0.5T_c$, $0.7T_c$, and $0.9T_c$ in decreasing order of gradient at the origin for the configurations of (a) **aa** and (b) **ab**. Here, $I_0 = S\Delta_B(0)k_F^2/4\pi\hbar$ with S being the interface area.




FIG. 2: Josephson current-phase relations of *BAB* junctions at $0.1T_c$ for $L = 0, \xi_0, 2\xi_0, 3\xi_0, 5\xi_0$, and $1000\xi_0$, in decreasing order of gradient at the origin for the configurations of (a) **aa** and (b) **ab**.

the gaps obey the BCS temperature dependence. We can observe current-phase curves with 4π rather than 2π periodicity and find some peak structures at lower temperatures. In particular, the structures become pronounced as the temperature decreases. In the Andreev picture, the Josephson current is carried by both a discrete part (Andreev bound states) and a continuous part of the excitation spectrum. This suggests that the peak structures and the 4π periodicity come from the Andreev bound states and the continuous spectrum, respectively.

It is known that Andreev bound states are localized near an interface. In the system with two interfaces that we consider here, the interaction between bound states near the two interfaces results in the current depending on the distance between the interfaces [23, 24]. In order to clarify the bound state contribution to the current, we investigate the current-phase relations of junctions with different *A*-phase widths at low temperatures. Figure 2 clearly shows that the peak structures are pronounced, while the 4π component is suppressed as L decreases. When $L \gg \xi_0$, the Andreev bound states near the two interfaces are isolated and their overlap is lost. A *BAB* junction is regarded as a series of *BA* and *AB* junctions. Since the total phase difference through the *BAB* junc-

tion is φ , the BA (AB) junction has $\varphi/2$ dependence, resulting the 4π periodicity in the current-phase characteristics. In addition, the current-phase relations in (a) and (b) merge into a single curve in the limit $L \rightarrow \infty$ because the current contributions from isolated interfaces are the same. When $L \rightarrow 0$, our model is reduced to the pinhole model in ^3He - B under a sufficiently large magnetic field applied parallel to the wall [10]. The surface free energy $-(\mathbf{H} \cdot R\hat{\mathbf{z}})^2$ produced by the applied magnetic field $\mathbf{H}$ is minimized if $\hat{\mathbf{n}}$ aligns in one of the four directions, $\mathbf{a}$, $\mathbf{b}$, $\mathbf{c}$, or $\mathbf{d}$. Therefore, $\hat{\mathbf{l}}$ plays a similar role to $\mathbf{H}$ in determining $\hat{\mathbf{n}}$ directions. This could be regarded as being a “pseudo-magnetic interface effect” that is inherent in BAB junction systems. The difference between $\mathbf{H}$ and the AB interface appears in the number of allowed $\hat{\mathbf{n}}$ directions: all four directions for $\mathbf{H}$, while only two directions in each $\hat{\mathbf{d}} = \pm\hat{\mathbf{l}}$ for the AB interfaces. Thus the curve for $L = 0$ in Fig. 2(b) is simply Yip’s π state [10], that is produced by the current cancellation due to quasiparticles with different $\hat{\mathbf{n}}$ -dependent effective phases. The curve in Fig. 2(a) corresponds to the curve for “AA” in Fig. 4 of Ref. 10. From the above discussion, the curve for “AC” cannot exist in a BAB junction. For $L \sim \xi_0$, the Andreev bound states near two interfaces maintain their overlap and the BAB junction behaves like a single junction. The current-phase relation still maintains the characteristics of the $L = 0$ curve.

Now let us revisit Fig. 1. The bound states localized near each interface overlap because $L \sim \xi_0$. Since the current contribution due to the Andreev bound states is dominant at low temperatures, 2π -periodic current-phase characteristics appear in the curves as peak structures. As the temperature becomes higher, the continuous spectrum contribution becomes large and obscures the Andreev bound state contribution. Therefore, we conclude that the peak structures appearing in Fig. 1 come from the Andreev bound states and the 4π periodicity is brought about by the continuous spectrum.

Throughout the above studies of BAB junctions, the current-phase characteristics reflected the configurations of textures and were modified by arranging the textures of each phase. Since our formula applies to any type of Josephson junction, systematic studies of current-phase relations reveal the internal structures due to spin-orbit degrees of freedom as well as their pairing symmetries.

In summary, we derived a general formula for Josephson current based on the Andreev picture. This formula made it possible to analyze any type of Josephson junction between unitary states with any symmetry, and was applied to novel BAB junctions in superfluid ^3He , which are accessible to experiments. It has been predicted that there are two types of current-phase relations associated with AB interfaces inherent in the system, acting sim-

ilarly to magnetic field $\mathbf{H}$ through $\hat{\mathbf{l}}$ in the A phase (“pseudo-magnetic interface effect”).

We thank E. V. Thuneberg, J. Viljas, G. E. Volovik, M. Krusius, L. Skrvák, J. Kopu, and D. Meacock for stimulating discussions. We are grateful to Y. Tanaka and Y. Ohashi for valuable input. We also thank Y. Asano for sending us his preprint. N. H. is indebted to H. Takayanagi and M. Paalanen for their continuous encouragement. This work was supported in part by the Waseda University Grant for Special Research Projects (2000A-875), by the COE Program “Molecular Nano-Engineering” from the Ministry of Education, Science and Culture, Japan, and by the Helsinki University of Technology and the National Technology Agency Tekes.

Note added.— During the preparation of this Letter, we learned of a theory [25] related to Josephson current in a similar system.

-
- [1] B. D. Josephson, *Adv. Phys.* **14**, 419 (1965).
 - [2] M. Sigrist and T. M. Rice, *J. Phys. Soc. Jpn.* **61**, 4283 (1992).
 - [3] D. A. Wollman *et al.*, *Phys. Rev. Lett.* **71**, 2134 (1993).
 - [4] R. Jin *et al.*, *Phys. Rev. B* **59**, 4433 (1999).
 - [5] M. Yamashiro, Y. Tanaka, and S. Kashiwaya, *J. Phys. Soc. Jpn.* **67**, 3364 (1998).
 - [6] C. Honerkamp and M. Sigrist, *Prog. Theor. Phys.* **100**, 53 (1998).
 - [7] S. Backhaus *et al.*, *Nature* **392**, 687 (1998).
 - [8] A. Marchenkov *et al.*, *Phys. Rev. Lett.* **83**, 3860 (1999).
 - [9] O. Avenel, Y. Mukharsky, and E. Varoquaux, *Physica B* **280**, 130 (2000).
 - [10] S.-K. Yip, *Phys. Rev. Lett.* **83**, 3864 (1999).
 - [11] J. K. Viljas and E. V. Thuneberg, *Phys. Rev. Lett.* **83**, 3868 (1999).
 - [12] A. F. Andreev, *Zh. Eksp. Teor. Fiz.* **46**, 1823 (1964) [*Sov. Phys. JETP* **19**, 1228 (1964)].
 - [13] A. Furusaki and M. Tsukada, *Solid State Commun.* **78**, 299 (1991).
 - [14] S. Yip, *Phys. Rev. B* **32**, 2915 (1985).
 - [15] J. Kurkijärvi, *Phys. Rev. B* **38**, 11184 (1988).
 - [16] I. O. Kulik and A. N. Omel’yanchuk, *Fiz. Nizk. Temp.* **3**, 945 (1977) [*Sov. J. Low Temp. Phys.* **3**, 459 (1977)].
 - [17] A. Furusaki and M. Tsukada, *Phys. Rev. B* **43**, 10164 (1991).
 - [18] Y. Tanaka and S. Kashiwaya, *Phys. Rev. B* **53**, R11957 (1996).
 - [19] D. J. Cousins *et al.*, *Phys. Rev. Lett.* **77**, 5245 (1996).
 - [20] R. W. Simmonds *et al.*, *Phys. Rev. Lett.* **84**, 6062 (2000).
 - [21] S. Yip, *Phys. Rev. B* **35**, 8733 (1987).
 - [22] E. V. Thuneberg, *Physica B* **178**, 168 (1992).
 - [23] M. Hurd and G. Wendin, *Phys. Rev. B* **51**, 3754 (1995).
 - [24] A. Nakayama, T. Furukawa, and Y. Okabe, *J. Appl. Phys.* **88**, 6605 (2000).
 - [25] Y. Asano, preprint.